



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

other parts forwarded to the National Museum by Mr. R. S. Allen for identification.

I have a cutting from a Richmond, Virginia, paper, name and date unknown, which notes the occurrence in New Kent County of four Wood Ibises, one of which was shot by Mr. W. J. Taylor and the head and wing exhibited in Richmond. The date was evidently the summer of 1893. Two, an adult and a young, were killed in Washington, July 2, 1892, as previously mentioned in 'The Auk,' (X, 1893, p. 91).—WILLIAM PALMER, *Washington, D. C.*

A North American Snipe New to the A. O. U. List.—Swainson in Sw. and Rich., *Fn. Bor.-Am.*, II, 1831, p. 501, describes *Scolopax leucurus* sp. n., from Hudson Bay, and says that "a single specimen of this bird in fine and perfect plumage, exists in the British Museum." It is a large species with 16 rectrices, whereof the three outer pairs are almost entirely white. Swainson's type is still extant in the British Museum, and has been identified with *Gallinago major* by Dr. Sharpe, who cites *S. leucurus* among the synonyms of this species, and catalogues Swainson's type from "Hudson's Bay" (*Cat. B. Brit. Mus.*, XXIV, 1896, p. 628 and p. 631). The specimen appears as one of a large lot acquired by the British Museum by purchase from the Hudson Bay Company many years ago, to none of which does any doubt of locality attach; and as there is obviously no reason to call Dr. Sharpe's identification in question, we may rely upon the record for the following addition to our Check-List:

[230.1.] ***Gallinago major* (Gm.).**

Greater Snipe.

Scolopax major GM. *Syst. Nat.* 1, 1788, p. 661.

Gallinago major KOCH, *Syst. Baier.* Orn. 1816, p. 313.

GEOG. DIST.—Europe, Asia and Africa. Accidental in North America ("Hudson Bay").

There may be a question about the specific name of this bird, under our Procrustean rules. According to Dr. Sharpe's synonymy, it is *Scolopax media* Frisch, 1763, and *Gallinago media* Gerini, 1773—both antedating Gmelin's *S. major*. Dr. Sharpe sensibly passes over *media* and adopts *major*—for to call a bird *media* in Latin and Greater or Double Snipe in English would be absurd. I hope some way can be found to square common sense with the Code in this case—if it cannot be, so much the worse for the latter.—ELLIOTT COUES, *Washington, D. C.*

The Occurrence of *Tryngites subruficollis* in the New England States.—There are several instances of the capture of the Buff-breasted Sandpiper in Connecticut and Massachusetts which do not appear to have been recorded. This Sandpiper, although common in certain sections in the West, is not very often met with along the North Atlantic seaboard.